1	
2	
3	Global warming shifts Pacific tropical cyclone location
4	
5	
6	
7	Tim L ¹ Min He Kwen ¹ Ming $Zhee^2$ long Seeng Kwe ³ ling lie Lue ⁴ and Weideng
/	Thin Li, Miniho Kwoli, Ming Zhao, Jong-Seolig Kug, Jing-Jia Luo, and weldolig
8	Yu ³
9	
10	¹ International Pacific Research Center, University of Hawaii, Honolulu, Hawaii
11	² GFDL, NOAA, Princeton, New Jersey
12	⁵ Korea Ocean Research and Development Institute, Korea
13	⁴ Research Institute for Climate Change, JAMSTEC, Japan
14	⁵ First Institute of Oceanography, State Oceanic Administration, China
15	
16	
17	
18	
19	
20	
21	
22	
23	
24	
25	
26	
27	
28	
29	
30	
31	
32 22	Corresponding Author:
33 24	Corresponding Author:
34 25	Tim Li International Pacific Research Center University of Heweii et Mance 1690
33 26	Fast Wast Dd Hanalulu HI 06822 E mail: timl:@hawaii adu
0C 27	Last- west Ru., Honolulu, FI, 90622. E-Illall. Ullill@llawall.edu
51	

Abstract

A global high-resolution (~ 40km) atmospheric general circulation model (ECHAM5 40 T319) is used to investigate the change of tropical cyclone frequency in the North 41 Pacific under global warming. A time slice method is used in which sea surface 42 temperature fields derived from a lower-resolution coupled model run under the 20C3M 43 (in which historical greenhouse gases in 20th century were prescribed as a radiative 44 forcing) and A1B (in which carbon dioxide concentration was increased 1% each year 45 from 2000 to 2070 and then was kept constant) scenarios are specified as the lower 46 boundary conditions to simulate the current and the future warming climate, 47 respectively. A significant shift is found in the location of tropical cyclones from the 48 western to central Pacific. The shift to more tropical cyclones in the central and less in 49 the western Pacific is not attributable to a change in atmospheric static stability, but to a 50 51 change in the variance of tropical synoptic-scale perturbations associated with a change 52 in the background vertical wind shear and boundary layer divergence.

54 **1. Introduction**

Tropical cyclones (TC) are among the most devastating weather phenomena 55 that can affect human life and economy. How global warming will affect TC activity is a 56 hotly debated topic [Webster et al., 2005; Emanuel, 2005; Landsea et al., 2006]. It has 57 been long recognized that TC genesis depends on sea surface temperature (SST) 58 because a higher SST provides TCs with high ocean thermal energy. In addition to SST, 59 TC genesis also depends on other dynamic and thermodynamic conditions such as 60 atmospheric static stability, humidity, perturbation strength, and vertical wind shear 61 [Gray, 1979; Wu and Lau, 1992; Emanuel and Nolan, 2004]. Particularly in the western 62 North Pacific, atmospheric circulation patterns rather than local SST play an important 63 role in interannual and interdecadal timescales [Chan, 2000; Matsuura et al., 2003; Ho 64 et al., 2004] and future projection [Yokoi and Takayabu, 2009]. 65

TCs originate from tropical disturbances. Statistically, only a small percentage 66 of the tropical disturbances eventually develop into TCs. With the increase of the global 67 SST and surface moisture, it is anticipated that more TCs would develop. However, 68 many climate models simulate a global decreasing trend of TC frequency [Sugi et al., 69 2002; McDonald et al., 2005; Hasegawa and Emori, 2005; Yoshimura et al., 2006; 70 Oouchi et al., 2006; Bengtsson et al., 2007]. One explanation of the decrease of TC 71 72 frequency is attributed to an increase of atmospheric static stability. This is because the 73 global warming leads to a larger increase of air temperature in the upper troposphere 74 than in the lower troposphere; as a result, the atmosphere becomes more stable, which suppresses the TC frequency [Sugi et al., 2002; Bengtsson et al., 2007]. If this is true 75 and it dominates the regional change of TC frequency, then one would expect the 76

decrease of TC frequency throughout all ocean basins. However, as shown by this study,

78

there are opposite trends of TC frequency between the western and central Pacific.

79 A concept of a relative SST warming was introduced recently to explain the change of basin-wide TC frequency under global warming [e.g., Swanson, 2008; Vecchi 80 et al., 2008; Knutson et al., 2008; Zhao et al. 2009, 2010]. In particular, Zhao et al. 81 [2009] found that both the present-day inter-annual variability of Atlantic hurricane 82 frequency and the inter-model spread in their simulated frequency response of 83 84 hurricanes to 21st century global warming projections (based on the IPCC AR4 models) can be well explained by a simple relative SST index defined as the Atlantic Main 85 Development Region SST minus the global tropical mean SST. While this result 86 suggests that the large-scale dynamic and thermodynamic conditions that are relevant to 87 TC genesis are closely tied to spatial distributions of SST in the Atlantic and eastern 88 Pacific [Zhao et al., 2010], such a simple relative SST index fails in explaining the 89 90 change in the western Pacific. Furthermore, it is worth mentioning that the relative SST 91 concept implies the importance of changes in atmospheric circulation in response to 92 change of SST distribution in affecting future TC frequency.

This study investigates the cause of shift of TC genesis locations in the Pacific in a warmer climate based on a high-resolution atmospheric general circulation model (AGCM). In the following, we first introduce the model and numerical experiments, and then we discuss the model results, with a focus on the shift of TC location and associated characteristics of mean state change. Finally, we summarize the main finding of this study.

99 2. Methodology and model description

100	AGCM used in this study is ECHAM5 [Roeckner et al., 2003] at a horizontal
101	resolution of T319 (about 40-km grid). This high-resolution global model is run at
102	Japan's Earth Simulator. SST, the lower boundary condition of the model, is derived
103	from a lower-resolution (T63) coupled version of the model (ECHAM5/MPI-OM)
104	[Jungclaus et al., 2006], which participated in the fourth assessment report of
105	intergovernmental panel for climate change (IPCC-AR4). Two different climate change
106	scenarios (20C3M and A1B) were applied. In 20C3M scenario, increasing historical
107	greenhouse gases in 20 th century were prescribed as a radiative forcing. In A1B scenario
108	carbon dioxide concentration was increased at a rate 1% per year till it reached 720 ppm
109	and was then kept constant. A 'time-slice' method [Bengtsson et al., 1996] was applied,
110	in which the high-resolution AGCM is forced by SST during two 20-year periods
111	(1980-1999 and 2080-2099). The two periods are hereafter referred to as 20C and 21C,
112	respectively.

Following Thorncroft and Hodges [2001], TCs in the model are determined based on the following three criteria: 1) 850-hPa vorticity is greater than $1.75 \times 10^{-6} \text{ s}^{-1}$, 2) warm core strength (represented by the difference between 850 and 250 hPa vorticity) exceeds $0.8 \times 10^{-6} \text{ s}^{-1}$, and 3) duration time exceeds 2 days. The selection of the parameter values is based on the least square fitting of the observed TC number in northern hemisphere in 20C.

119 **3. Results**

Figure 1 shows the geographical distribution of TC genesis locations in the Pacific in the 20C and 21C simulations. In 20C, TCs form primarily over the western and eastern Pacific, similar to the distribution of the observed genesis locations [Gray,

123 1979]. In 21C, however, more TCs shift their genesis locations to the Central Pacific. As
124 seen from the difference map (Fig. 1c), there are two notable TC decrease and increase
125 regions over the Pacific. One is over the North western Pacific (NWP) and the other the
126 North central Pacific (NCP). The numbers of TCs in 21C decreases by 31% over NWP
127 but increases by 65% over NCP. Thus the high-resolution AGCM simulations illustrate
128 two opposite TC trends in NWP and NCP.

To demonstrate that the result above is not a specific feature of the ECHAM5 129 130 model and its projected SST warming pattern, we also examined the simulation results from the GFDL global 50-km resolution AGCM (HiRAM2.1, for detailed model 131 description and its realistic simulation of the annual, inter-annual and decadal variations 132 of TC activity, readers are referred to Zhao et al. 2009) with an ensemble SST warming 133 pattern from 18 IPCC AR4 models [Zhao et al., 2009]. The two models have different 134 treatments in various physical parameterization schemes. For example, for convective 135 136 heating parameterization ECHAM5 uses a mass flux scheme while HiRAM2.1 uses a 137 modified Bretherton (2004) scheme. Figure 1d shows the difference in TC genesis number between 21C and 20C from the GFDL model. An opposite TC trend can also be 138 139 seen between NWP and NCP, supporting the ECHAM5 result.

To understand the cause of this distinctive shift in TC location with global warming, we diagnose in the following the dynamic and thermodynamic conditions in northern summer (July – October), when a majority of TCs occur, over the NWP (5-25°N, 110°E-160°E) and NCP (5-25°N, 180-130°W) regions, respectively. First we examine the change of atmospheric stability in both the regions. Figures 2a and 2b show the vertical profiles of the atmospheric potential and equivalent potential temperatures

averaged over NWP and NCP. The upper-level air temperature increases at a greater rate 146 than that at lower levels in both the regions. As the static stability is measured by the 147 vertical gradient of the potential temperature, the result implies that the atmosphere 148 149 becomes more stable under the global warming in both the regions. Due to the effect of atmospheric humidity (which increases more in the lower troposphere), the vertical 150 gradient of equivalent potential temperature approximately remains same from 20C to 151 21C. This implies that the atmospheric convective instability remains unchanged in both 152 the regions. Thus, both the static stability and the convective instability changes cannot 153 154 explain the opposite trends of TC frequency between NWP and NCP.

A further analysis reveals that the fundamental cause of the opposite TC trends 155 lies in the change of dynamic conditions. As we know, TCs originate from the tropical 156 disturbances such as synoptic wave trains and easterly waves [e.g., Lau and Lau, 1990]. 157 The 21C simulation shows an increased variability of synoptic-scale disturbances over 158 159 the NCP region but a decreased synoptic activity over the NWP region. Here the 160 strength of the synoptic-scale disturbances is represented by the variance of the 850-hPa 161 vorticity field that is filtered at a 2-8 day band using Lanzcos digital filter [Duchon, 162 1979]. The difference map (Fig. 2c) shows a remarkable decrease of the synoptic-scale variance over (10-20N, 110-140E) but an increase of the variance in (10-20N, 180-163 130W). The importance of synoptic activity on future projection of TC genesis 164 frequency was also pointed out by Yokoi and Takayabu [2009]. Thus, the decreasing 165 trend in NWP is caused by the reduced synoptic-scale activity whereas the increasing 166 trend in NCP is caused by the strengthening of synoptic disturbances. 167

168

We argue here that the contrast of the synoptic-scale activity between NWP and

NCP results from the change of the background vertical wind shear (Fig. 3a) and low-169 level divergence (Fig. 3c). It has been pointed out that the easterly shear and low-level 170 convergence of the background mean flow favor the development of tropical 171 172 disturbances [Wang and Xie, 1996; Li, 2006; Sooraj et al., 2008]. In the 20C simulation, the western Pacific and Indian monsoon regions have a prevailing easterly wind shear in 173 association with large-scale convective heating. This easterly wind shear becomes 174 weaker in the warming climate (Fig. 3a). The weakening of the easterly wind shear 175 corresponds to a weakened western Pacific monsoon, as seen from Fig. 3b. The 176 177 weakening of the easterly shear, along with low-level divergence to its east (Fig. 3c), further suppresses the development of tropical disturbances in NWP. In contrast, the 178 179 easterly wind shear and low-level convergence in NCP are strengthened, and so is the 180 precipitation (Fig. 3). They favor the development of TCs in NCP.

What causes the change of the vertical shear and the low-level divergence? We 181 182 argue that the change of the background vertical shear and the low-level divergence is 183 closely related to changes of the trade wind in the tropics (Fig. 3c). Many IPCC-AR4 models predict an El Nino-like warming pattern in the Pacific under the global warming 184 185 climate state [Solomon et al., 2007], that is, a greater SST warming occurs in the tropical eastern and central Pacific compared to that in the tropical western Pacific. As a 186 187 result, the zonal SST gradient is reduced across the tropical Pacific. The reduced zonal SST gradient decreases the trade wind [Lindzen and Nigam, 1987] and weakens the 188 Walker circulation. Furthermore, GCM simulations with a uniform SST warming also 189 190 produce a weakening of the Walker circulation, which can be understood in terms of the energy and mass balance of the ascending branch of the circulation [Held and Soden, 191 2006; Vecchi et al., 2006]. Therefore, both the reduced zonal SST gradient and the 192

193 global mean SST warming decrease the trade wind.

The weakening of the trades leads to, on the one hand, the decrease of the boundary-layer convergence in the western Pacific monsoon region and, on the other hand, the increase of the boundary convergence in NCP (Fig. 3c). The former suppresses the monsoon convective heating and decreases the easterly shear in NWP, whereas the latter strengthens the local boundary layer moisture convergence and convective heating and thus increases the easterly wind shear in NCP.

200 4. Summary and discussion

The shift of TC location in the Pacific due to the global warming is investigated using a high-resolution (~ 40km) global AGCM (ECHAM5). It is noted that TC genesis number decreases significantly in the western Pacific but increases remarkably in the central tropical Pacific. Such a feature is also seen in the GFDL HiRAM2.1 forced with an IPCC-AR4 ensemble SST warming pattern.

The cause of the two opposite TC trends is investigated. As the atmospheric static stability increases in both the regions under global warming, this stability control mechanism cannot explain the increasing trend over the central Pacific. The major factor that accounts for the distinctive opposite TC trends lies in the dynamic condition in the atmosphere. It is the change of strength of synoptic disturbances that is primarily responsible for the distinctive changes in the TC frequency between the western and central Pacific.

213 Physical processes responsible for the distinctive TC frequency changes in the
214 warming climate are discussed below. The global warming weakens the trade winds in

the Pacific through both the effect of the uniform warming and the decrease of zonal 215 216 SST gradients across the tropical Pacific. In the western Pacific, the weakened Walker 217 circulation causes the anomalous boundary layer divergence that leads to a weakening 218 of the western North Pacific monsoon rainfall. The weakening of the monsoon heating further causes a decrease in the background easterly shear. The decreased easterly shear 219 220 and associated weakening of the low-level convergence to its east reduce the synoptic activity and thus decrease the TC genesis frequency. In contrast, in the central Pacific, 221 222 an anomalous boundary layer convergence is induced due to the reduced SST gradient 223 and higher SST. This enhances the atmospheric convection and induces the easterly 224 shear in situ. The easterly shear and the low-level convergence favor the growth of the 225 synoptic disturbances, resulting in a TC frequency increase in the future warming 226 climate.

Projection on future cyclogenesis frequency is subject to a wide range of uncertainties including uncertainties in model physics and SST warming patterns [Knutson et al., 2010]. ECHAM5 tends to overestimate TC genesis frequency in the central Pacific in 20C. Thus a caution is needed to interpret the results. The shift of projected TC activity in the Pacific may pose a great threat to millions' people living in Hawaii and central Pacific islands.

233

Acknowledgments. This work was supported by ONR grants N000140810256 and N000141010774 and NRL grant N00173-09-1-G008 and by the International Pacific Research Center that is sponsored by the Japan Agency for Marine-Earth Science and Technology (JAMSTEC), NASA (NNX07AG53G) and NOAA (NA17RJ1230). This is 238 SOEST contribution number 8016 and IPRC contribution number 720.

References

241	Bengtsson, L., M. Botzet, and M. Esch (1996), Will greenhouse gas-induced warming
242	over the next 50 years lead to higher frequency and greater intensity of hurricanes?
243	<i>Tellus</i> , 48A , 57-73.
244	Bengtsson, L., K. I. Hodges, M. Esch, N. Keenlyside, L. Komblush, J. J. Luo, and T.
245	Yamagata (2007), How many tropical cyclones change in a warmer climate. Tellus,
246	59A , 539-561.
247	Bretherton, C.S., J.R. McCaa, and H. Grenier, 2004: A new parameterization for shallow
248	cumulus convection and its application to marine subtropic cloud-topped
249	boundary layers. Part I: description and 1D results. Mon. Wea. Rev., 132, 864-882.
250	Chan, J. C. L., 2000: Tropical cyclone activity over the Western North Pacific
251	associated with El Niño and La Niña events. J. Climate, 13, 2960-2972.
252	Duchon, C. E. (1979), Lanczos filtering in one and two dimensions, J. Appl. Meteoro.,
253	18 , 1016-1022.
254	Emanuel, K. A., and D. S. Nolan (2004), Tropical cyclone activity and global climate.
255	In: Proc. of 26 th Conference on Hurricanes and Tropical Meteorology, pp. 240-
256	241., American Meteorological Society, Miami, FL.
257	Emanuel, K.A. (2005), Increasing destructiveness of tropical cyclones over the past 30

436,	686-688
	436,

259	Gray, W. M. (1979), Hurricanes: Their formation, structure and likely role in the tropical
260	circulation. Meteorology over the Tropical Oceans, D.B. Shaw (Ed.), Royal
261	Meteorological Society, 155–218.
262	Hasegawa, A., and S. Emori (2005), Tropical cyclones and associated precipitation over
263	the western North Pacific: T106 atmospheric GCM simulation for present-
264	day and doubled CO2 climates. Scientific Online Letters on the Atmosphere, 1, 14
265	5–148.
266	Held, I. M. and B. J. Soden, (2006), Robust responses of the hydrological cycle to
267	global warming. J. Climate, 19, 56865699.
268	Ho, CH., JJ. Baik, JH. Kim, DY. Gong, and CH. Sui, 2004: Interdecadal changes
269	in summertime typhoon tracks. J. Climate, 17, 1767-1776
270	Jungclaus, H., and coauthors (2006), Ocean circulation and tropical variability in the co
271	upled model ECHAM5/MPO-OM. J. Clim., 19, 3952-3972.
272	Knutson, T., J. Sirutis, S. Garner, G.A Vecchi and I.M.Held (2008), Simulated
273	reduction in Atlantic hurricane frequency under twenty-first-century warming
274	condition. Nature Geoscience, doi:10.1038/ngeo202.

275 Landsea, C. W., B. A. Harper, K. Hoarau, and J. A. Knaff (2006), Can we detect trends

276	in extreme tropical cyclones? Science, 313, 452-454.
277	Lau, KH., and NC. Lau (1990), Observed structure and propagation characteristics of
278	tropical summertime synoptic-scale disturbances. Mon. Wea. Rev., 118, 1888-1913.
279	Li, T. (2006), Origin of the Summertime Synoptic-Scale Wave Train in the Western
280	North Pacific. J. Atmos. Sci., 63, 1093-1102.
281	Lindzen, R.S., and S. Nigam (1987), On the role of sea surface temperature gradients in
282	forcing low level winds and convergence in the tropics. J. Atmos. Sci., 44, 2440-
283	2458.
284	Matsuura, T., M. Yumoto, and S. Iizuka, 2003: A mechanism of interdecadal variability
285	of tropical cyclone activity over the western North Pacific. Clim. Dyn., 21, 105-
286	117.
287	McDonald, R. E., and coauthors (2005), Tropical storms: Representation and diagnosis
288	in climate models and the impacts of climate change. Clim. Dyn., 25, 19–36.
289	Oouchi, K., and coauthors (2006), Tropical cyclone climatology in a global-warming
290	climate as simulated in a 20km-mesh global atmospheric model: Frequency and
291	wind intensity analyses. J. Meteorol. Soc. Japan, 84, 259–276.
292	Roeckner, E., and coauthors, 2003: The atmospheric general circulation model
293	ECHAM5. Part I: Model description. Max Planck Institute for Meteorology

294 Rep. 349, 127 pp.	
-----------------------	--

295	Solomon, S, and coauthors (eds) Climate Change 2007: The Physical Science Basis
296	(Cambridge University Press for the Intergovernmental Panel on Climate Change,
297	2007).
298	Sooraj, K. P., D. Kim, JS. Kug, SW. Yeh, F-F. Jin, and IS. Kang (2008), Effects of
299	the low frequency zonal wind variation on the high-frequency atmospheric
300	variability over the tropics. Clim. Dyn., 33 , doi 10.1007/s00382-008-0483-6 (CA).
301	Sugi, M., A. Noda, and N. Sato (2002), Influence of the global warming on tropical
302	cyclone climatology: An experiment with the JMA global model. J. Meteorol. Soc.
303	Japan, 80 , 249–272.
304	Swanson, K. L., (2008), Nonlocality of Atlantic tropical cyclone intensities.
305	Geochem. Geophys. Geosyst., 9, Q04V01, doi:10.1029/2007GC001844.
306	Thorncroft, C., and K. Hodges (2001), African easterly wave variability and its
307	relationship to Atlantic tropical cyclone activity. J. Clim., 14, 1166–1179.
308	Vecchi, G. A., K. Swanson, and B. Soden (2008), Whither hurricane activity.
309	Science, 322, 687-689.
310	Vecchi, G. A., B. J. Soden, A. T. Wittenberg, I. M. Held, A. Leetmaa and M. J.
311	Harrison (2006), Weakening of tropical Pacific atmospheric circulation due

312	to anthropogenic forcing. Nature 441, 73-76.
313	Wang, B., and X. Xie (1996), Low-frequency equatorial waves in sheared zonal flow.
314	Part I: Stable waves. J. Atmos. Sci., 53, 449-467. (1996).
315	Webster, P. J., G. J. Holland, J. A. Curry, and HR. Chang (2005), Changes in tropical
316	cyclone number, duration, and intensity in a warming environment. Science, 309,
317	1844-1846.
318	Wu, G., and NC. Lau (1992), A GCM simulation of the relationship between tropical-
319	storm formation and ENSO. Mon. Wea. Rev., 120, 958-977.
320	Yokoi, S., and Y. N. Takayabu, 2009: Multi-model projection of global warming impact
321	on tropical cyclone genesis frequency over the Western North Pacific. J. Meteor.
322	Soc. Japan, 525-538.
323	Yoshimura, J., M. Sugi, and A. Noda (2006), Influence of greenhouse warming on
324	tropical cyclone frequency. J. Meteorol. Soc. Japan, 84, 405-428.
325	Zhao, M., I. M. Held, SJ. Lin and G. A.Vecchi (2009), Simulations of global
326	hurricane climatology, interannual variability, and response to global
327	warming using a 50km resolution GCM. J. Climate, 22, 6653-6678.
328	Zhao, M., I. M. Held and G. A. Vecchi (2010), Retrospective forecasts of the
329	hurricane season using a global atmospheric model assuming persistence of

330 SST anomalies. Mon. Wea. Rev., in press.

332 Figure Legends

333	Figure 1 TC genesis number at each 2.5°x2.5° box for a 20-year period derived from
334	T319 ECHAM5 for (a) 20C, (b) 21C, and (c) difference between (b) and (a) (21C-
335	20C). In (c) red (orange) shaded areas indicate 95% (90%) confidence level. (d)
336	same as (c) except from the GFDL HiRAM2.1 model with an ensemble
337	SST pattern averaged from 18 IPCC-AR4 models (dashed line denotes the
338	95% confidence level).
339	Figure 2 Vertical profiles of potential and equivalent potential temperatures (unit: K) at
340	20C and 21C averaged over (a) NWP and (b) NCP and the variance difference (c)
341	of synoptic-scale (2-8-day) vorticity at 850 hPa (unit: 10^{-10} s ⁻²) in northern summer
342	(July-October) between 21C and 20C, with shaded areas indicating a 95%
343	confidence level or above (with an F test).
344	Figure 3 Difference (21C – 20C) fields of (a) the vertical shear of zonal wind (200 hPa
345	minus 850 hPa, unit: m s ⁻¹), (b) precipitation (unit: mm day ⁻¹), and (c) velocity
346	potential (unit: s ⁻¹) and wind (unit: m s ⁻¹) at 850 hPa during northern summer
347	(July-October). Areas that exceed the 95% confidence level (Student's t test) are
348	shaded (for contour) and plotted (for vector).
349	







Figure 3